

*"I am stunned at the amazing results I've been getting with L.A.S.T.!"*

*"L.A.S.T. is an invaluable set of techniques that have improved outcomes for my patients!"*

*"It was a great course...my patient said it was the best treatment he's ever had!"*




**L.A.S.T.**  
Ligamentous Articular Strain Technique

# Canadian Massage Conference 2015

## Techniques for the Shoulder

Presented by Robert Libbey, RMT



# Canadian Massage Conference 2015 Shoulder Course Manual

L.A.S.T. Ligamentous Articular Strain Technique®™ 2010-2015

No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without either the prior permission of the publisher or license permitting restricted copying in Canada issued by the Canadian Intellectual Property Office, Place du Portage I, 50 Victoria St., Room C-114 Gatineau, Quebec K1A 0C9

## Note:

Knowledge and best practice in this field are constantly changing. As new research and experience broaden our knowledge, changes in practice and treatment may become necessary or appropriate. Readers are advised to check the most current information concerning the frequency, intensity, duration, precautions, indications, and contraindications concerning treatment of patients and their conditions. It is the responsibility of the practitioner, relying on their own experience and knowledge of the patient, to make diagnoses, to determine frequency, intensity and duration and the best treatment for each individual patient, and to take all appropriate safety precautions. To the fullest extent of the law, the author does not assume any liability for any injury and or damage.

Book design by Robert Libbey Massage Therapist Corporation and Original 72 Creative

Cover design by Robert Libbey Massage Therapist Corporation and Original 72 Creative

Photos by Original 72 Creative

Editing assistance from Leslie Ste. Marie

Copyrighted 2015



**“IN AN**

**ASYMMETRICAL**

**RAENIL-NON**

**FSESDTBEMCK,**

**SMALL** CHANGES

**CAN RESULT IN**

**DISPROPORTIONATELY**

**LARGE**

**EFFECTS”**



# Contents

LIGAMENOUS ARTICULAR STRAIN TECHNIQUES GLOBAL PAIN THERAPIES .....	4
BREATHING .....	9
BREATHING TECHNIQUES.....	10
CLINICAL PRINCIPLES .....	12
THE APPLICATION OF L.A.S.T. ....	13
POST TREATMENT.....	14
<b>SHOULDER TECHNIQUES.....</b>	<b>15</b>
Anterior Cervical Fascia.....	16
Anterior Sternoclavicular Ligament.....	17
Posterior Sternoclavicular Ligament .....	18
Costoclavicular Ligament/Upper Mediastinum .....	19
Subclavious .....	20
Corocoid Ligaments.....	21
Pectoralis Minor, Coracobrachialis, .....	22
Short Head Biceps .....	22
Long Head Biceps/Glenohumeral Capsule .....	23
Teres Major/Minor .....	24
GH Capsule Disengagement .....	25
Respiratory Diaphragm.....	28
COMMON MISTAKES/ DO'S AND DON'TS .....	29
LIGAMENT PAIN REFERRAL PATTERN POSTERS .....	31
LIGAMENOUS ARTICULAR STRAIN TECHNIQUE COURSES.....	32
REFERENCES.....	33
BIBLIOGRAPHY.....	34
THE CLINICAL OBSERVATION OF MUSCLE ENERGY TECHNIQUES AND LIGAMENOUS ARTICULAR STRAIN IN 2 CASES OF CERVICAL DISC HERNIATION WITH THORACIC OUTLET SYNDROME.....	37



# Techniques for the Shoulder

## Workshop Agenda:

### 9am-5pm

9:00am	Registration, Introduction & Scientific Theory for L.A.S.T.
10:00am	Shoulder Relevant Anatomy, CIs, Indications, Precautions, Pathologies, Kinesiology 1. Anterior Cervical Fascia 2. Ant./Post. Sternoclavicular Ligaments
10:30am	Student Exchange
11:30am	Shoulder Relevant Anatomy, CIs, Indications, Precautions, Pathologies, Kinesiology 3. Costoclavicular Ligament and Upper Mediastinum 4. Subclavious 5. Corocoid Ligaments
12:00am	Student Exchange
1:00pm	Lunch
2:00pm	Shoulder Relevant Anatomy, CIs, Indications, Precautions, Pathologies, Kinesiology 6. Pectoralis Minor, Coracobrachialis, Short Head Biceps 7. Long Head Biceps 8. Teres Major/Minor
2:30pm	Student Exchange
3:30pm	Shoulder and Global Effects On It Relevant Anatomy, CIs, Indications, Precautions, Pathologies, Kinesiology 9. GH Capsule Disengagement 10. Respiratory Diaphragm
4:00pm	Student Exchange
5:00pm	End of day



## Forward Thinking

The information presented in this manual before you, is an opportunity to advance and update the original context first put forth by A.T. Still. This is a wholly different era where evidence informed techniques are sought after in an effort to advance the manual therapy profession to it's highest potential.

Presented here are the same timeless principles viewed through a modern lens. They are applied with the mindset of providing a simple, precise, principle based, evidence informed technique that can be incorporated into techniques already being utilized by the manual therapist.

At the time of writing this, after 20 years of practice and 16 years of research (both historically and scientifically informed), self development and education has lead me to this historically innovative manual therapy technique.

Although the techniques described in this manual seem separate, they are connected through the ability of the therapist to look locally and see systemically. We are an asymmetrical nonlinear feed back system. Everything is connected.

In the book "Movement, Stability & Lumbopelvic Pain", Andry Vleeming uses the term "*Ligamentous Stocking*" to describe the connectedness of fibrous soft-tissue structures of the lumbar vertebrae to the sacrum. In 2009, Jaap van der Wal stated that there is a joint stability system, in which muscular tissue and RDCT (regular dense (collagenous) connective tissue) interweave and function mainly in an "*in series*" situation rather than an "*in parallel*" situation. Thus, in vivo, the periarticular connective tissue is loaded and stretched both by the movement of related skeletal parts and by the tension of the muscle tissue inserting to this connective tissue. Ligaments are considered RDCT's.

I put it to you that this "*Ligamentous Stocking*" is organism wide, connecting not just the vertebral column to the sacrum, but also connecting the various appendicular interdigitations of membranous, capsular, ligamentous and periosteal fibrous tissues to the axial fibrous tissues.

The trend today is to treat separate tissue from other separate tissue with a disconnect from the whole organism. We treat carpal tunnel at the wrist with out looking systemically to find that the wrist was only the last in a chain of events and compensations for something that happened months years, decades ago in a completely different area.



## Canadian Massage Conference 2015 Shoulder Course Manual

Daily we see patients who seem to have been treated by everyone, everywhere, and some how have come to us. We are in most cases the “end of the road” before surgery or they come to us decades after surgery. Their physical body is screaming out information that many do not stop to hear or see. You can see it in their eyes.

We still utilize the ancient laying of hands to help someone in pain and discomfort. We are here to help, to serve, to listen with our hands, eyes, ears, heart, soul and intuition. We are here to connect with another organism.

Going forward, use today's evidence informed science to help you and your patient understand possibly what physiological process is currently occurring; but use your common sense, intuition and most importantly guidance from your patient's physiology to dictate the rate, course and direction of the treatment.

Manual therapy is an ART and a science. Treat the organism rather than the segment. Change your perspective to treating densities and temperatures. Aim for “*ever-changing balance*” in an asymmetrical nonlinear feedback system.

Keep Forward Thinking!



## LIGAMENTOUS ARTICULAR STRAIN TECHNIQUES

### Global Pain Therapies

#### Ligamentous Articular Strain Techniques - A manual treatment approach for ligamentous articular injuries and for the whole body

Robert Libbey, RMT

#### ABSTRACT

*The scope of practice for the treatment of the ligamentous, capsular and fascial tissues by Manual Therapists (Massage Therapy) limits the therapist to utilizing their hands along with active patient participation. There is a wide range of research documenting the types of mechanoreceptors and their locations within the ligamentous/capsular tissues. A lesser-understood ligamento-muscular reflex has also been documented with limited research. The specific mechanoreceptors and ligamento-muscular reflex are responsive to manual stimulation techniques within the scope of practice for manual therapy. Utilizing specific manual techniques can have positive effects in attenuating the development of scar tissue, adhesions, inflammatory processes and proprioceptive and kinesthetic neurological deficits. Combining these specific manual techniques with therapies such as Prolotherapy minimizes the long-term effects from ligamentous articular injuries and accelerates the recovery period for patients.*

*Journal of Prolotherapy, 2012;4:e886-e890.*

**KEYWORDS:** ACL, capsular tissue, fascial tissue, L.A.S.T., ligamentous tissue, massage therapy, Prolotherapy.

“It’s not perverted function but a wrong environment that results in the distorted appearance of function. Function is always true to its environment. Function is dependent upon its environment. Therefore, any change in any part of the environment that is not in tune or balance, will distort the function of the matter so involved.”

- Thomas Schooley, DO<sup>1</sup>

There is a wealth of information concerning the treatment of ligamentous, capsular and fascial tissues within the scope of practice of Medicine, Physiotherapy and Naturopathic Medicine. In the Massage Therapy (Manual Therapy) profession where our scope of practice only allows us to treat utilize our hands along with active patient participation; there is very little information or research.

In this article, I discuss a technique developed in the early 1900’s to which I have applied today’s current research. When combined with therapies such as Prolotherapy, this technique has the potential to accelerate the recovery period for patients suffering from ligamentous articular injuries.

The ligamentous system is part of the fascial tissue referred to as the **Multimicrovascular Collagenic Dynamic Absorbing System**. The role of this rubbery elastic shock absorbing system, found everywhere in our body, is to avoid reaching a threshold of resistance at which the collagen might shear resulting in injury.<sup>2</sup>

Ruptured or injured ligaments are a source of mechanical problems. These injuries cause deficiencies in fine and gross motor controls, lost, altered or impaired sensation, various types of pain/discomfort, joint laxity and can lead to articular surface injury. This dysfunctional state eventually leads to an increased risk of additional injury to surrounding capsules, tendons, neurovasculature and other nearby tissue, can cause a change in sensation thresholds, impaired sensation and impaired reflex muscular activation.<sup>3</sup>

If left untreated or if treated inefficiently, chronic inflammation develops and permanent disability of the affected tissues results. Patients suffer from continual pain, inflammation, neurological implications, muscle stiffness and instability of affected joints, deficient sensation and sensory perception, impaired motor performance, deficient synergy muscle activation, spasms and hyper excitability of the muscular system.<sup>3</sup>





LIGAMENTOUS ARTICULAR STRAIN TECHNIQUE

MECHANORECEPTORS

Ligaments have significant input to sensation and contribute to the synergistic activation of muscles. Afferent mechanoreceptors are found everywhere throughout our connective tissues. They are responsible for kinesthetic and proprioceptive sensation.<sup>3</sup>

The four types of these sensory nerve endings or mechanoreceptors that manual therapists have influence on are:

- Golgi Receptors
- Ruffini Receptors
- Pacini Receptors
- Interstitial Receptors

We will briefly look at these four receptors and the effects manual treatment has on them and the body as a whole. (See Table 1.)<sup>4, 5, 6, 7, 8, 9, 10, 11</sup>

In a study presented at the Third International Fascia Research Congress, Viklund et al. concluded that specific myofascial receptor techniques might have a longer lasting effect than classical (Swedish) massage techniques. They suggested that “therapists might be encouraged to aim their soft tissue techniques to a lesser area where there is known to be high density of mechanoreceptors”.<sup>12</sup>

Manual techniques that target these mechanoreceptors have been proven to affect both the local blood supply and the local tissue viscosity.<sup>11</sup> Research is now confirming the common clinical finding that slow, deep tissue techniques have both local and systemic effects.

LIGAMENTO-MUSCULAR REFLEX

Ligaments are not passive tissue. Ligaments are highly dynamic and non-stationary predictable organs. Afferent mechanoreceptors in ligaments trigger a ligamento-muscular reflex activation of associated muscles. It has been suggested that a reflex might exist between these sensory receptors in the ligaments and surrounding associated muscles. This Ligamento-muscular reflex may directly or indirectly alter the load inflicted on to a ligament.<sup>3</sup>

Table 1. The four types of sensory nerve endings or mechanoreceptors and the effects manual treatment has on them and the body as a whole.	
<b>Receptor:</b>	<b>Golgi</b>
<b>Location:</b>	<ul style="list-style-type: none"> <li>• all dense connective tissues</li> <li>• peripheral joints ligaments</li> <li>• joint capsules</li> <li>• myotendinous junctions</li> <li>• attachment areas of aponeurosis</li> </ul>
<b>Responds to:</b>	slow stretch techniques
<b>Results in:</b>	a decrease in active muscle tone
<b>Receptor:</b>	<b>Ruffini</b>
<b>Location:</b>	<ul style="list-style-type: none"> <li>• tissues associated with regular stretching</li> <li>• outer layer of joint capsules dura mater</li> <li>• peripheral joints ligaments</li> <li>• muscle fasciae</li> <li>• the deep dorsal fascia of the hand</li> </ul>
<b>Responds to:</b>	constant, slow and deep pressure slow shear forces
<b>Results in:</b>	a lowering of sympathetic nervous system activity
<b>Receptor:</b>	<b>Pacini</b>
<b>Location:</b>	<ul style="list-style-type: none"> <li>• deep portions of joint capsules</li> <li>• deeper spinal ligaments</li> <li>• investing muscular fasciae of antebrachial, crural,</li> <li>• abdominal fasciae, masseter, lateral thigh, plantar and palmar tissues &amp; peritoneum</li> </ul>
<b>Responds to:</b>	<ul style="list-style-type: none"> <li>• rapid changes in pressure</li> <li>• vibratory/oscillatory techniques</li> <li>• HVLA's</li> </ul>
<b>Results in:</b>	an increase local proprioceptive attention and self-regulation
<b>Receptor:</b>	<b>Free/Interstitial Nerve Endings</b>
<b>Location:</b>	<ul style="list-style-type: none"> <li>• all fascial tissue</li> <li>• periosteum</li> <li>• interosseous membranes</li> <li>• tendons</li> <li>• fascial connections to bones</li> </ul>
<b>Responds to:</b>	<ul style="list-style-type: none"> <li>• deep, slow or steady manual pressure</li> <li>• Treatment to periosteum, interosseous membranes, and fasciae connected to bones</li> </ul>
<b>Results in:</b>	<ul style="list-style-type: none"> <li>• autonomic functions: changes in heart rate, blood pressure, respiration, increase or decrease blood pressure, sensation of position and movement, increased vagal activity, global muscle relaxation, alterations in local fluid dynamics and tissue metabolism</li> <li>• plasma extravasation</li> </ul>

The Ligamento-muscular reflex has been shown to exist in most joints of the extremity and in the spine. Muscular activity elicited by this reflex allows muscles and ligaments to work together as a unit in maintaining joint stability. This reflex may play a role in the preservation of joint stability, inhibiting muscles that destabilize the joint or increasing antagonist co-activation to stabilize the joint.<sup>3</sup>



## LIGAMENTOUS ARTICULAR STRAIN TECHNIQUE

For many years we learned in our anatomy classes that ligaments and muscles were separate entities. Even many of our most popular anatomy texts still portray ligaments and muscles as separate tissues running from one bone to another, not connected to any surrounding tissues. In reality though, these and the surrounding tissues are inseparable. In 2009, van der Wal published a paper in which he determined that ligaments are mostly arranged in series with the muscles, not parallel.<sup>13</sup> When you contract a muscle, the ligaments are automatically engaged, assisting in the stabilization of a joint, no matter what its position, during both concentric and eccentric contractions.<sup>14</sup>

## INJURIES

A state of hypertonicity is created in muscles when a ligament has been injured, or replaced. For example, ACL ruptures and replacements can cause one or all of the quadriceps, hamstring and gastrocnemius muscles to become hypertonic as a result of a lack of the ligamento-muscular reflex. This hypertonic state is also due to the removal of the mechanoreceptors of the original ACL. A large percentage of the receptors are located near the ends of the ligaments as they attach into the sub-chondral layer. This remnant is often removed in order to attach the substituted tissue. Although the stability of the joint has been re-established, patients regularly complain of not only having an increase in tonicity of the muscle groups, but also a deficiency in both proprioception and kinaesthetic awareness. The manual therapist is then charged with the task of attempting various treatment techniques to reduce this hypertonicity. Their efforts are rarely successful and both the patient and therapist become frustrated and discouraged. Thus begins the circle of referral from one therapist to another.<sup>5, 15, 16, 17, 18</sup>

## HISTORY OF L.A.S.T.

Ligamentous Articular Strain Technique (L.A.S.T.) was developed to be primarily an Indirect Osteopathic technique. The technique is based on principles and techniques developed by Andrew Taylor Still, DO. The majority of the techniques initially developed were called traction methods, known as "Indirect Techniques." Several of these techniques were Ligamentous Articular Strain Techniques and some of them also formed the basis of myofascial release techniques. Many of the techniques in the early 20th century were modified by therapists that wanted to focus on a quicker, more direct method

of treating. These "Direct Techniques," became known primarily as high velocity-low amplitude techniques.<sup>1, 19</sup>

The author has re-defined the techniques to suite the Scope of Practice for Massage Therapists. "Indirect Techniques" follow the permitted motions of the dysfunctional tissues into the direction of ease. "Direct Techniques" match the reciprocal tension of the dysfunctional tissues taking the tissues only to the first tissue resistance barrier.

## USING L.A.S.T. IN TREATMENT

L.A.S.T. affects the connective tissues of the body, mainly ligaments, joint capsules, fascia, muscles, tendons and indirectly, lymphatic and blood flow and the autonomic nervous system.<sup>11, 20</sup>

L.A.S.T. is a principle-based technique. The basic principles consist of disengaging the dysfunctional tissues from their protective position. The practitioner follows the affected tissues into a position of exaggeration of the injury. This exaggeration leads to a state where all movement within the affected tissues halts. A release of tension within the tissue is felt, allowing the ligaments to draw the articulations back into a more normal balanced relationship.<sup>19</sup>

After an assessment of the affected articular structures, the practitioner palpates for areas of dysfunction or an increased densification within the injured tissues. Slowly disengaging the tissues, (direct or indirect techniques) the therapist exaggerates the permitted motions of the tissues to their end barrier. The reciprocal tension is a tension that is developed between the practitioner loading into the tissue and the tissue reciprocally loading back into the practitioner. The therapist now waits at this point of tissue exaggeration until he or she feels the ligaments draw the articulations back into a normal balanced position. A re-assessment of the permitted motion in the tissues should confirm more suppleness and mobility along with an improvement in kinaesthetic and proprioceptive awareness. (See Figures 1-4.)

## L.A.S.T., TGF-B1 AND THEIR POTENTIAL ROLE IN PREVENTING EXCESSIVE SCAR TISSUE DEVELOPMENT

Solomonow states that the acute inflammation in ligaments sets in within several hours may last several weeks and up to 12 months! Only up to 70% recovery



LIGAMENTOUS ARTICULAR STRAIN TECHNIQUE



Figure 1. Hand placement for treatment of Annular Ligament.



Figure 3. Hand placement for treatment of ASIS attachment for Inguinal Ligament.



Figure 2. Sidelying position for treatment of Iliolumbar ligament.



Figure 4. Hand placement on patient for treatment of C1 dysfunction.

has been documented. Chronic inflammation can build up over several weeks, months or years depending on dose-duration levels. Rest and recovery of as long as two years only allows partial recovery, full recovery has never been reported.<sup>3</sup>

Bouffard et al. published a study documenting the effects of brief static tissue stretch on TGF- $\beta$ 1. TGF- $\beta$ 1 plays a key role in connective tissue regulating the response of fibroblasts to injury, remodelling, scarring, and pathological production of fibrosis.<sup>23</sup> Langevin et al. stated that in cases of minor sprains and repetitive motion injuries, scarring is mostly detrimental since it can contribute to maintaining the chronicity of tissue stiffness, abnormal movement patterns, and pain. Reducing scar and adhesion formation using stretch and mobilization is especially important for internal tissue injuries and inflammation involving fascia and organs.<sup>21</sup>

The results of the Bouffard and Langevin studies showed that brief, moderate amplitude (20–30% strain) stretching of connective tissue decreases both TGF- $\beta$ 1 and collagen synthesis.

Langevin et al. proposed that therapies which briefly stretch tissues beyond the habitual range of motion (eg, massage) locally inhibit new collagen formation for several days after, and thus, prevent and/or ameliorate soft tissue adhesions.<sup>6,7,19,23,24</sup>

Thomas Schooley, DO stated, “Function is dependent upon its environment.” I believe that by utilizing the principles of L.A.S.T in various stages of injury, practitioners have the potential to profoundly affect the course of tissue healing. By changing the neuromuscular physiological environment and decreasing the SNS response to injury, a response of decreased TGF- $\beta$ 1

could lead to decrease in fibrosis and decrease in fascial stiffening of the surrounding and injured tissues. This approach could overall positively affect the functionality of the patient.

#### MY EXPERIENCE WITH L.A.S.T.

In our office, we treat a variety of injuries ranging from acute to chronic. These injuries lead to the symptoms described earlier in this article. Optimally, I prefer to treat in the acute stages of an injury. By attempting to create as optimal an environment for tissue healing as possible, patients have reported being less symptomatic. Combining L.A.S.T with Prolotherapy treatment, administered by our Naturopathic Physician, creates an opportunity to prevent chronic issues from occurring thereby improving the quality of life for the patient. ■

#### REFERENCES:

1. The Osteopathic Cranial Association. (1953). Journal of the Osteopathic Cranial Association.
2. Endo Vivo (Producer), & Guimberteau, J. C. (Director). (2004). *Strolling under the skin* [DVD].
3. Solomonow M. Ligaments: a source of musculoskeletal disorders. *Journal of Bodywork and Movement Therapies*, 2009;13(2): 136-54. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/19329050>.
4. Cottingham JT. (1985) *Healing Through Touch*. Boulder, CO: Rolf Institute Publications.
5. Krosgaard et al. Cruciate ligament reflexes. *J Electromyogr Kinesiol*. 2002 Jun;12(3):177-82. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/12086811>.
6. Langevin HM, et al. Mechanical signaling through connective tissue: A mechanism for the therapeutic effect of acupuncture. *FASEB J*. 2001;15:2275-2282.
7. Langevin HM, et al. Dynamic fibroblast cytoskeletal response to subcutaneous tissue stretch ex vivo and in vivo. *Am J Physiol Cell Physiol*. 2005;288:C747-C756.
8. Sakada S. (1974). Mechanoreceptors in fascia, periosteum and periodontal ligament. *Bull Tokyo Med Dent Univ*, 21 (Suppl.), 11-13.
9. Stilwell D. (1957). Regional variations in the innervation of deep fasciae and aponeuroses. *The Anatomical Record*, 127(4), 635-653.
10. van den Berg, F. & Cabri, J. (1999). *Angewandte Physiologie – Das Bindegewebe des Bewegungsapparates verstehen und beeinflussen*. Stuttgart, Germany: Georg Thieme Verlag
11. Schleip R. *Dynamic Body: Exploring Human Form, Expanding Human Function* Fascia as a Sensory Organ: A Target of Myofascial Manipulation.
12. Viklund P. (2012) Comparison of ankle joint dorsiflexion after classical massage or specific myofascial receptor massage technique on the calf muscle. Third International Fascia Research Congress: Basic Science and Implications for Conventional and Complementary Health Care. Munich, Germany: Elsevier GmbH.
13. van der Wal J. The architecture of the connective tissue in the musculoskeletal system—an often overlooked functional parameter as to proprioception in the locomotor apparatus. *International Journal of Therapeutic Massage and Bodywork*. 2009 Dec;2(4).
14. Myers T. (2011). Dynamic Ligaments: Re-visioning the Fascia as a Body-Wide Regulatory System. *Massage Magazine*.
15. Dhillon MS, et al. Proprioception in anterior cruciate ligament deficient knees and its relevance in anterior cruciate ligament reconstruction. *Indian J Orthop*. 2011 Jul;45(4):294-300. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/21772620>.
16. Lee BI, et al. Immunohistochemical study of mechanoreceptors in the tibial remnant of the ruptured anterior cruciate ligament in human knees. *Knee Surg Sports Traumatol Arthrosc*. 2009 Sep;17(9):1095-101. Epub 2009 Jun 16. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/19533097>.
17. Melnyk M, et al. Changes in stretch reflex excitability are related to “giving way” symptoms in patients with anterior cruciate ligament rupture. *J Neurophysiol*. 2007 Jan;97(1):474-80. Epub 2006 Aug 30. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/16943314>.
18. Swanik CB, et al. Reactive muscle firing of anterior cruciate ligament-injured females during functional activities. *J Athl Train*. 1999 Apr;34(2):121-9. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/16558554>.
19. Speece CA, et al. (2009). *Ligamentous Articular Strain: Osteopathic Manipulative Techniques for the Body* (Revised edition). Seattle, WA: Eastland Press.
20. Coote JH, et al. The response of some sympathetic neurons to volleys in various afferent nerves. *The Journal of Physiology*. 1970;208(02): 261-278.
21. Langevin HM, et al. Pathophysiological model for chronic low back pain integrating connective tissue and nervous system mechanisms. *Med Hypotheses*. 2007;68:74-80.
22. Bouffard NA, et al. Tissue stretch decreases soluble TGF B1 and Type-1 pro-collagen in mouse subcutaneous connective tissue: evidence from ex vivo and in vivo models. *Journal of Cellular Physiology*. 2008;214: 389-395, 2008.
23. Langevin HM, et al. Evidence of connective tissue involvement in acupuncture. *FASEB J*. 2002;16:872-874.
24. Langevin HM, et al. Subcutaneous tissue fibroblast cytoskeletal remodeling induced by acupuncture: evidence for a mechanotransduction-based mechanism. *J Cell Physiol*. 2006;207:767-774.



## Breathing

Utilizing breathing as a technique when performing manual techniques is greatly underutilized by a majority of today's therapists. Therapists that utilize concepts from techniques such as CST are familiar with incorporating breath into their treatments; but why not more of us.

Evidence supports a link between breathing difficulties and back pain. Since trunk muscles perform both postural and breathing functions, disruption in one function can negatively impact the other.

Altered breathing mechanics changes respiratory chemistry causing smooth muscle constriction, altered electrolyte balance and decreased tissue oxygenation. These changes can profoundly impact any body system. Increased excitability in the muscular and nervous systems may be most relevant to a manual therapist. (McLaughlin L, et al., Breathing evaluation and retraining as an adjunct to manual therapy, *Manual Therapy* (2010), doi:10.1016/j.math.2010.08.006 )

Breathing is unquestionably a key function of the human body; it sustains life by providing oxygen needed for metabolism and removing the by-product of these reactions, carbon dioxide. Breathing, however, has other functions apart from the ventilation of air and the maintenance of oxygen and carbon dioxide. Breathing affects motor control and postural stability and plays several roles in physiological and psychological regulation. Breathing can influence homeostatic functions in other systems including the autonomic nervous system, the circulatory system, chemical regulation and metabolism.

Breathing becomes dysfunctional when the person is unable to breathe efficiently or when breathing is inappropriate, unhelpful or inefficient in responding to environmental conditions and the changing needs of the individual.

Impairment of the functions of breathing affects people's lives, challenging homeostasis, creating symptoms and compromising health. The efficiency with which breathing fulfills its various functions can be diminished because of musculo-skeletal dysfunction, disease, chronic psychological stress or other factors that affect respiratory drive and respiratory control. The neurological control of breathing shows high levels of neuroplasticity as shown by its ability to adapt to a wide range of internal and external conditions.

Breathing therapy generally aims to either correct dysfunctions of breathing or enhance its functions. Breathing, unlike most physiological functions, can be controlled voluntarily and it can serve as an entry point for physiological and psychological regulation. (International Journal of Osteopathic Medicine Volume 12, Issue 3 Pages 78-85, September 2009)

Quite simply, with diminished function of breath, our physiological systems cease to function at their maximum potential. As this process becomes chronic, the physiological environment becomes more acidic, functioning in a more sympathetic, fight, flight state slowing the recuperative processes of our immune system.

There are an abundance of instructional books, videos and classes educating us on the importance of breath and its physiological effects on all our systems.

Simply put, without it, we cease.

## Breathing Techniques

There are various opinions on how, when and where to breath. As always, L.A.S.T. primarily looks to change the environment that the dysfunctional tissues are attempting to function within. There are 2 ways that we change the environment with LAST, holding of inhalation or exhalation by the patient.

### Inhalation Instruction:

Once you have disengaged and exaggerated the injured tissue(s) from their dysfunctional holding position, instruct the patient to breath in as deep as possible utilizing as much of their lungs and diaphragm as they can. Once they reach the top of the breath, they are to hold this breath as long as comfortably possible before exhaling. This process can be repeated if necessary to create the effected change.

### Exhalation Instruction:

Once you have disengaged and exaggerated the injured tissue(s) from their dysfunctional holding position, instruct the patient to breath out as deep as possible. Once they have comfortably forced all the air out of their body to their fullest extent, they are to hold this as long as comfortably possible before inhaling. This process can be repeated if necessary to create the effected change.

“The quieter the mind  
The stiller the hands  
The less movement we make  
The more we are able to perceive involuntary movement”  
- James Jealous, DO

“It’s not perverted function but a wrong environment that results in the distorted appearance of function. Function is always true to its environment. Function is dependent upon its environment. Therefore, any change in any part of the environment that is not in tune or balance will apparently distort the function of the matter so involved.”

- Thomas Schooley, DO



## Clinical Principles

### Contraindications

acute fracture, acute instabilities - dislocations, immediate acute sprain/strains, acute periosteal tears, acute RA, infection, acute systemic conditions

### Precautions

general health of the patient, stages of healing post-injury, medications, recent injectable medications, plates/pins, degeneration of various stages, fractures, various degrees of sprains/strains, fusions, instabilities, periosteal tears

### Indications

decreased AROM/PROM, pain, tension, stiffness, discomfort, acute/sub-acute/chronic sprain/strains, carpal tunnel, numbness, tingling, vascular changes

### Direct and Indirect Techniques

Through manual manipulation, we can perceive changes in patients' tissues that originate in the afferent mechanoreceptors within the tissue. By gently, yet firmly loading into the specific tissue, we feel for the reciprocal tension of the tissue pressing back into our hands. Instead of forcing through the barrier of tissue directing a release as we see fit, we come to just before the barrier and allow the tissue to release in the direction of ease (Indirect Technique).

In some circumstances, however, the tissue requires that we match the reciprocal tension in the tissue and wait for the balance to occur (Direct Technique).

When do we use one or the other? The tissue determines the course of treatment and in many cases; both indirect and direct techniques are utilized in order to bring balance to the tissue/system.



## The Application of L.A.S.T.

L.A.S.T. is a principle-based technique: Disengage, Exaggerate, and Balance.

Find the reciprocal tension and motion permitted within the membranes.

*Disengage* the tissue (using direct or indirect techniques, or a combination of both) through all planes of motion, not on any particular axis, within a joint in the direction of ease. This is known as an *Exaggeration*.

Hold the position until you come to a point of suspension: a neutral position or *Balance*. Wait in that position until a softening of the tissues occurs.

Once the tissue has completed balancing, re-assess the motion permitted within the joint.



)

# Post Treatment

No change

Better

Worse

No better

No worse

Better later

Worse later



# Shoulder Techniques

















# Pectoralis Minor, Coracobrachialis, Short Head Biceps

**Patient position:**

Supine on a table with affected shoulder off the table

**Therapist position:**

Standing at the head table



**Technique:**

Caution - If the shoulder has been previously dislocated, position the arm on the table for support

This technique is similar for all of these tissues; the only differences are in the direction of force and hand placement.

Pectoralis minor muscle: Locate the attachment at the coracoid process. Matching the reciprocal tension of the tissues, maintain steady, balanced pressure in the direction of it's attachments onto ribs 3-5, until you feel a release of the tension.

Coracobrachialis muscle/short head of the biceps: Place your thumbs just lateral and inferior to the coracoid process applying the same steady, balanced pressure in a lateral and posterior direction. Matching the reciprocal tension of the tissues, maintain steady, balanced pressure until you feel a release of the tension.

---

---

---

---

---

---

---

---

---

---

## Long Head Biceps/Glenohumeral Capsule

**Patient position:**

Supine on a table with affected shoulder off the table



**Therapist position:**

Seated at the head table



**Technique:**

Caution - If the shoulder has been previously dislocated, do not perform this technique!

This technique targets the longhead of the biceps and it's synovial sheath as it travels inside the capsule attaching onto the supraglenoid tubercle of the scapula. It also assists in challenging the tissues of the glenoid capsule as the head of the humerus moves anteriorly and inferiorly.

The normal range of motion of the glenohumeral capsule into extension is 45-60 degrees. Depending on the severity/chronicity of injury, these ranges may not be able to attain in one treatment, therefore gaining this range should be structured over a period of treatments.

First position the arm off the table at a point that is comfortable for the patient. Gently extend the elbow and pronate the wrist. Slowly start to bring the whole arm into further extension. Stop as the patients discomfort starts to increase. Instruct the patient to perform the inhalation breathing technique. Once the patient fully exhales, proceed to bring the arm into more extension once again.

---

---

---

---


---



# GH Capsule Disengagement

**Patient position:**  
Supine on the table

**Therapist position:**  
Seated at the head of the table, next to the patient on the same side to be treated



**Technique:**

WARNING RISK OF HARM: Do not attempt this technique with a patient suffering from a previous GH Dislocation(s)!

This technique challenges the tissues that connect the inferior aspect of the AC joint to the superior aspect of the GH joint. It encourages the head of the humerus anteriorly and inferiorly, challenging the GH joint capsule along with its attachments to the GH Fossa, Labrum and AC joint.

Release the Teres muscles before performing this technique.

Place the thenar web of one hand under the posterior GH joint to encourage the GH capsule anteriorly and inferiorly.

With your other hand, gently contact the patients flexed elbow. The intent is to slowly drawing the arm into a position of abduction and external rotation. With both hands working together as one unit, compress both hands towards each other through the humerus, adjusting the direction of forces. Feel the maximum reciprocal tension in the tissues between your two hands. Maintain steady, balanced pressure until you feel a release of the tension occur.

---

---

---

---

---

---


---




---

---

---

## Respiratory Diaphragm









## Common Mistakes/ Do's and Don'ts

### Don't

Do not think that this is a cure.

We are just changing the environment of the body to increase it's functioning.

"It's not perverted function but a wrong environment that results in the distorted appearance of function. Function is always true to its environment. Function is dependent upon its environment. Therefore, any change in any part of the environment that is not in tune or balance will apparently distort the function of the matter so involved."

- Thomas Schooley, DO

### Do

Remember your principles:

Disengage, Exaggerate, Balance

Acute, Sub-acute, Chronic

Listen and follow the tissues/body

### Don't

Over treat your patients with this technique tomorrow.

Risk of harm

Can be too much change for the patient in one session

### Don't

Rush through the barrier.

Direct and Indirect Techniques

### Do

Listen and feel for the RECIPROCAL TENSION in the tissues.



## Don't

Work quickly.

## Do

Slow down!

“The quieter the mind

The stiller the hands

The less movement we make

The more we are able to perceive involuntary movement”

- James Jealous, DO

## Don't

Treat on hardware you have never seen before  
get x-rays, CT/MRI reports

## Don't

Think you will be proficient in this technique immediately!

It takes time and Patients! LOTS of patients - 500-1000!

## Do you feel Overwhelmed?


### 1<sup>st</sup> Thing to do!

Incorporate 1-2 techniques/patient/treatment to start.

Don't over treat - It can be too much for the patient to handle!



## Ligament Pain Referral Pattern Posters



These visually stunning, full-color posters are an invaluable diagnostic and educational tool for you and your patients.

Over the last 18yrs I've recognized a distinct, immediate and ongoing problem. My patients were complaining of pain referral patterns that didn't seem to match up with the standard trigger points, dermatome or sclerotome patterns.

For years I researched scientific journals and resources all the while continually charting the referral patterns described to me by my patients.

This extensive work has culminated in my creation of this set of LIGAMENT PAIN REFERRAL PATTERN posters!

This stunning artwork is original and doesn't exist anywhere else!

These 2 posters graphically demonstrate over 40 ligament pain referral patterns for: Spine & Upper Extremity, Spine & Lower Extremity

Multiple referral patterns are displayed for each region of the body; Spine, Shoulder, Forearm & Hand, Hip & Pelvis, Knee, Leg & Foot.

If you have patients complaining of referred pain that you just can't figure out, if you treat joint dysfunction and want to explain more in depth the discomfort your patients are feeling, if you want to add more value to your practice and your patients... these posters will be an asset!

These visually stunning, full-color, high glossy charts are an invaluable diagnostic and educational tool for you and your patients.

Help your patients visually see the cause of their complaints!

To order now simply go to [www.lastsite.ca](http://www.lastsite.ca).

# Ligamentous Articular Strain Technique Courses

## Multi Day Courses

Upper Body & Extremities

Lower Body & Extremities

Thorax, Abdomen & Pelvis

Bowstring & Diaphragms

## 1 Day Courses

Postural Correction Techniques

Thorax Treatment Techniques

Treatment Techniques for Shoulder

Treatment Techniques for Hip & Pelvis

Treatment Techniques for Foot, Leg & Knee

## Online Courses

Techniques for Shoulder

Techniques for Leg & Foot

Techniques for Knee

Techniques for Hip & Pelvis

Techniques for Postural Correction

Techniques for Sternum

For course schedules, visit [www.lastsite.ca](http://www.lastsite.ca)



## References

- Coote, J. H. & Pérez-González, J. F. (1970). The response of some sympathetic neurons to volleys in various afferent nerves. *The Journal of Physiology*, 208(02), 261-278.
- Cottingham, J. T. (1985) *Healing through Touch*. Boulder, CO: Rolf Institute Publications.
- Mitchell, J. H. & Schmidt, R. F. (1977). Cardiovascular reflex control by afferent fibers from skeletal muscle receptors. *Handbook of Physiology* (eds. Shepherd JT et al.) Sect. 2, vol. III, part 2, 623-658. Bethesda, MD: American Physiological Society.
- Sakada, S. (1974). Mechanoreceptors in fascia, periosteum and periodontal ligament. *Bull Tokyo Med Dent Univ*, 21 (Suppl.), 11-13.
- Stilwell, D. (1957). Regional variations in the innervation of deep fasciae and aponeuroses. *The Anatomical Record*, 127(4), 635-653.
- van den Berg, F. & Cabri, J. (1999). *Angewandte Physiologie - Das Bindegewebe des Bewegungsapparates verstehen und beeinflussen*. Stuttgart, Germany: Georg Thieme Verlag.
- Yahia, L., Rhalmi, S., Newman, N. & Isler, M. (1992). Sensory innervation of human thoracolumbar fascia. *Acta Orthopaedica Scandinavica*, 63(2), 195-197.



## Bibliography

- Barral, J. P. (1991). *The Thorax*. Seattle, WA: Eastland Press.
- Barral, J. P. & Mercier, P. (2006). *Visceral Manipulation (Revised Edition)*. Seattle, WA: Eastland Press.
- Barral, J. P. (2007). *Visceral Manipulation II (Revised Edition)*. Seattle, WA: Eastland Press.
- Becker, R. E. (1997). *Life in Motion*. Portland, Oregon: Stillness Press LLC.
- Cantu, R. I. & Grodin, A. J. (2001) *Myofascial Manipulation, Theory and Clinical Application: Second Edition*. New York, NY: Aspen Publishers.
- Chu, D., LeBlanc, R., D'Ambrosia, P., D'Ambrosia, R., Baratta, R. V. & Solomonow, M. (2003). Neuromuscular Disorder in Response to Anterior Cruciate Ligament Creep. *Clinical Biomechanics*, 18(3), 222-30. Retrieved from [http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=12620785&ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=12620785&ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Diederichsen, L. P., et al. (2004). Reflexes in the Shoulder Muscles Elicited from the Human Coracoacromial Ligament. *Journal of Orthopedic Research*, 22(5), 976-83. Retrieved from [http://www.ncbi.nlm.nih.gov/pubmed/15304268?ordinalpos=24&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/pubmed/15304268?ordinalpos=24&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- DiGiovanna, E. L., Schiowitz, S. & Dowling, D. J. (2004). *An Osteopathic Approach to Diagnosis and Treatment: Third Edition*. Philadelphia, PA: Lippincott Williams & Wilkins.
- Dixon, M. (2006). *Joint Play the Right Way for the Axial Skeleton*. Vancouver, BC: Arthrokinetic Publishing.
- Eastland Press (Producer), & Barral, J. P. (Director). (2002). *Visceral Manipulation [DVD]*. Seattle, WA.
- Edmond, S. L. (1993). *Manipulation Mobilization Extremities and Spinal Techniques*. St. Louis, MO: Mosby.
- Endo Vivo (Producer), & Guimberteau, J. C. (Director). (2004). *Strolling under the skin [DVD]*.
- Endo Vivo (Producer), & Guimberteau, J. C. (Director). (2008). *The Skin Excursion [DVD]*.
- Endo Vivo (Producer), & Guimberteau, J. C. (Director). (2010). *Muscle Attitudes [DVD]*.
- Findley, T. W. & Schleip, R. (2007). *First International Fascia Research Congress: Basic Science and Implications for Conventional and Complementary Health Care*. Munich, Germany: Elsevier GmbH.
- Findley, T. W. et al. (2009). *Second International Fascia Research Congress: Basic Science and Implications for Conventional and Complementary Health Care*. Munich, Germany: Elsevier GmbH.
- Gorman, D. (2005). *The Body Moveable: Fifth Edition*. Guelph, ON: Learning Methods Publications.
- Hamwee, J. (2000). *Zero Balancing: Touching the Energy of Bone*. Berkley, CA: North Atlantic Books.
- Hoppenfeld, S. (1976). *Physical Examination of the Spine & Extremities*. Upper Saddle River, NJ: Prentice Hall.
- Integral Anatomy Productions, LLC (Producer), & Hedley, G. (Director). (2005). *The Integral Anatomy Series, Vol. 1: Skin and Superficial Fascia [DVD]*.
- Integral Anatomy Productions, LLC (Producer), & Hedley, G. (Director). (2006). *The Integral Anatomy Series, Vol. 2: Deep Fascia and Muscle [DVD]*.



- Integral Anatomy Productions, LLC (Producer), & Hedley, G. (Director). (2006). The Integral Anatomy Series, Vol. 3: Cranial and Visceral Fasciae [DVD].
- Integral Anatomy Productions, LLC (Producer), & Hedley, G. (Director). (2009). The Integral Anatomy Series, Vol. 4: Viscera and their Fasciae [DVD].
- Jealous, J. Perceptual Studies No. 1: The Sutherland Odyssey [CD]. United States: Direction of Ease, LLC.
- Jealous, J. Balanced Membranous Tension: No. 1 [CD]. United States: Direction of Ease, LLC.
- Jealous, J. Balanced Membranous Tension: No. 2: [CD]. United States: Direction of Ease, LLC.
- Jealous, J. The Patient's Neutral: No. 1 [CD]. United States: Direction of Ease, LLC.
- Jealous, J. (2006). Our Hands [CD]. United States: Direction of Ease, LLC.
- Johansson, H., Sjölander, P. & Sojka, P. (1991). A Sensory Role for the Cruciate Ligaments. *Clinical Orthopaedics and Related Research*, 268, 161-78. Retrieved from [http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=2060205&ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=2060205&ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Konishi, Y., Fukubayashi, T. & Takeshita, D. (2002). Mechanism of Quadriceps Femoris Muscle Weakness in Patients with Anterior Cruciate Ligament Reconstruction. *Scandinavian Journal of Medicine and Science in Sports*, 12(6), 371-5. Retrieved from [http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=12453165&ordinalpos=9&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=12453165&ordinalpos=9&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Konishi, Y., Fukubayashi, T. & Takeshita, D. (2002). Possible Mechanism of Quadriceps Femoris Weakness in Patients with Ruptured Anterior Cruciate Ligament. *Medicine and Science in Sports and Exercise*, 34(9), 1414-8. Retrieved from [http://www.ncbi.nlm.nih.gov/pubmed/12218732?ordinalpos=36&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/pubmed/12218732?ordinalpos=36&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Lederman, E. (2005). *The Science and Practice of Manual Therapy* (2<sup>nd</sup> ed.). Philadelphia, PA: Churchill Livingstone.
- Myers, T. (2011). Dynamic Ligaments: Re-visioning the Fascia as a Body-Wide Regulatory System. *Massage Magazine*.
- Netter, F. H. (1998). *Atlas of Human Anatomy*. Philadelphia, PA: Saunders Elsevier.
- Nicholas, A. S., Nicholas, E. A. (2008) *Atlas of Osteopathic Techniques* (1<sup>st</sup> ed.). Baltimore, MD: Lippincott, Williams & Wilkins.
- The Osteopathic Cranial Association. (1953). *Journal of the Osteopathic Cranial Association*.
- Park, H. B., Koh, M., Cho, S. H., Hutchinson, B. & Lee, B. (2005). Mapping the Rat Somatosensory Pathway from the Anterior Cruciate Ligament Nerve Endings to the Cerebrum. *Journal of Orthopaedic Research*, 23(6), 1419-24. Retrieved from [http://www.ncbi.nlm.nih.gov/pubmed/15921874?ordinalpos=17&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/pubmed/15921874?ordinalpos=17&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Phillips, D., Petrie, S., Solomonow, M., Zhou, B. H., Guanque, C. & D'Ambrosia, R. (1997). Ligamentomuscular Protective Reflex in the Elbow. *The Journal of Hand Surgery*, 22(3), 473-8. Retrieved from



- [http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=9195457&ordinalpos=26&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=9195457&ordinalpos=26&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Sakamoto, N., Yamashita, T., Takebayashi, T., Sekine, M. & Ishii, S. (2001). An Electrophysiologic Study of Mechanoreceptors in the Sacroiliac Joint and Adjacent Tissues. *Spine*, 26(20), E468-71. Retrieved from [http://www.ncbi.nlm.nih.gov/pubmed/11598526?ordinalpos=50&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/pubmed/11598526?ordinalpos=50&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Schleip, R. *Dynamic Body: Exploring Human Form, Expanding Human Function* Fascia as a Sensory Organ: A Target of Myofascial Manipulation.
- Schleip, R. (2004). 5<sup>th</sup> Interdisciplinary World Congress on Low Back & Pelvic Pain [oral presentation]. Melbourne, Australia.
- Schultz, R. L. & Freitis, R. (1996). *The Endless Web: Fascial Anatomy and Physical Reality*. Berkeley, CA: North Atlantic Books.
- Solomonow, M. (2003). Ligaments: A Source of Work-Related Musculoskeletal Disorders, [presentation at the 2003 STAR Symposium].
- Solomonow, M. (2006). Sensory-motor Control of Ligaments and Associated Neuromuscular Disorders. *Journal of Electromyography and Kinesiology*, 16(6), 549-67. Retrieved from [http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=17045488&ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed\\_ResultsPanel.Pubmed\\_RVDocSum](http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=17045488&ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum)
- Solomonow, M. (2009). Ligaments: A Source of Musculoskeletal Disorders. *Journal of Bodywork and Movement Therapies*, 13(2), 136-54. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19329050>
- Speece, C. A., Crow, W. T. (2009). *Ligamentous Articular Strain: Osteopathic Manipulative Techniques for the Body* (Revised edition). Seattle, WA: Eastland Press.
- Still, A. T. (1992). *Osteopathy: Research and Practice*. Seattle, WA: Eastland Press.
- Sutherland, W. G. (1998). *Contributions of Thought: The Collected Writings of William Garner Sutherland* (2<sup>nd</sup> ed.). Yakima, WA: Sutherland Cranial Teaching Foundation.
- van der Wal, J. (2009). The Architecture of the Connective Tissue in the Musculoskeletal System—An Often Overlooked Functional Parameter as to Proprioception in the Locomotor Apparatus. *International Journal of Therapeutic Massage and Bodywork*, 2(4).





# The clinical observation of muscle energy techniques and ligamentous articular strain in 2 cases of cervical disc herniation with thoracic outlet syndrome

International Journal of Osteopathic Medicine (2015) 18, 63–70



**International  
Journal of  
Osteopathic  
Medicine**

www.elsevier.com/ijos

## CASE REPORT

# The clinical observation of muscle energy techniques and ligamentous articular strain in 2 cases of cervical disc herniation with thoracic outlet syndrome



Min-yeong Kim, Eun Hya Chi, Jin-ho Lee, In-Hyuk Ha\*

*Jaseng Spine and Joint Research Institute, Jaseng Medical Foundation, 858 Eonju-ro, Gangnam-gu, Seoul, Republic of Korea*

Received 18 November 2013; revised 7 March 2014; accepted 16 October 2014

### KEYWORDS

Thoracic outlet syndrome;  
Cervical disc herniation;  
Muscle energy technique;  
Ligamentous articular strain

**Abstract** This case report presents two successful treatment outcomes of cervical disc herniation (CDH) with thoracic outlet syndrome (TOS) treated using manual therapy technique of the muscle energy technique (MET), ligamentous articular strain (LAS) and additional herbal medicine intake, acupuncture and pharmacopuncture sessions. Significant improvements were reported in the outcome measures at admission and at discharge for Visual Analogue Scale (VAS) and Neck Disability Index (NDI) scores and physical examinations followed by approximately three weeks of hospital admission. The successful outcomes suggest that integrative conservative management focused on muscle energy techniques (MET) and ligamentous articular strain (LAS) are effective in contributing to the conservative management of cervical disc herniation (CDH) with thoracic outlet syndrome (TOS). © 2014 Elsevier Ltd. All rights reserved.

## Introduction

Thoracic outlet syndrome (TOS) is a relatively rare condition prevalent in 8% of the general

population<sup>1,2</sup> whereas cervical disc herniation (CDH), a leading medical concern in the adult population,<sup>3–5</sup> has a prevalence of over 3.3 cases per 1000.<sup>6</sup> In CDH, neurological symptoms typically present concurrently with TOS,<sup>2</sup> making diagnosis difficult and demands both objective and subjective examinations.<sup>7–9</sup> CDH causes structural anomalies and medical imaging is often beneficial

\* Corresponding author. Tel.: +82 2 3218 2188; fax: +82 2 3218 2244.

E-mail address: hanihata@gmail.com (I.H. Ha).

<http://dx.doi.org/10.1016/j.ijosm.2014.10.008>  
1746-0689/© 2014 Elsevier Ltd. All rights reserved.



In establishing the diagnosis, TOS is defined as a complex condition involving the compression of either neurological or vascular elements passing through the thoracic outlet. The subclavian vein, artery and lower part of the brachial plexus are often involved<sup>10-13</sup> manifesting symptoms including pain, numbness and other related symptoms.<sup>10</sup> These symptoms are often aggravated by movement of the shoulder, cervical spine and head or by raising the upper limbs.<sup>10</sup>

Anatomically, the subclavian artery, subclavian vein and the lower part of brachial plexus travel through the thoracic outlet<sup>14</sup> and different clinical signs are observed depending on the affected site. Neurogenic TOS mostly presents with neurological symptoms, such as paraesthesia, weakness in the upper extremities, and pain. In cases where vascular elements are involved, symptoms of pallor, paraesthesia and coldness are often present due to arterial ischaemia in arterial TOS, and swollen upper extremities and cyanosis due to subclavian vein obstruction are observed in venous TOS.<sup>15</sup>

The overlapping signs and symptoms of CDH and TOS lead to issues with the differential diagnosis in

many cases, as well as issues with the selection of treatment approaches. No reports have been presented on the effect of Muscle Energy Technique (MET)<sup>16</sup> and LAS (Ligamentous Articular Strain)<sup>17</sup> for the treatment of CDH with associated TOS. The two cases in the present paper had previously undergone injection therapies, medications and other medical treatments for their CDH with associated TOS. Although no improvement was reported with these medical approaches, significant improvement was observed with manual therapy using MET and LAS. This case report presents two cases of the management of CDH with TOS using these two techniques.

## Clinical history

### Case 1

A 49-year-old female presented complaining of posterior cervical pain with right arm pain and numbness (Fig. 1) following an accident that




Fig. 1 Body chart – case 1.




Fig. 2 C-spine MRI: C3/4/5/6/7 HVD & stenosis.

involved bumping into a glass door which occurred 10 months prior to her initial visit. The cervical spine x-ray, CT and MRI scans revealed CDH with stenosis at C3/C4, C4/C5, C5/C6 and C6/C7 (Fig. 2). She was hospitalised at a local hospital for 3 days where she received traction therapy, injections, medications and nerve block injection therapy. Some improvements were observed initially but the pain remained. The patient was subsequently admitted to Jaseng Hospital of Korean Medicine<sup>36</sup> for 21 days. At admission, the patient reported a VAS score of 10 and a NDI score of 30. Cervical spine range of motion (ROM) testing revealed restrictions in right lateral bending of 20° (normal >45°). She complained that her pain was worse at night and indicated on the NDI that she was averaging 4 h of sleep per night. Positive findings in all three physical examinations were reported (Spurling test, Adson's test and Roos test). Cervical spine x-ray (Fig. 3) revealed a narrowing of the disc space, posterior marginal spurring and degenerative endplate change at C4/5, C5/6 and C6/7. At discharge, the patient reported a VAS score of 2 and a NDI score of 21. Normal c-spine ROM and negatives in all physical examinations were reported. At discharge, she reported

significant reduction in pain and her sleep was no longer affected. The c-spine ROM was reported as normal and all physical examinations reported negative.

### Case 2

A 35-year-old male patient presented with posterior cervical pain left arm numbness and pain (Fig. 4) which was worse at night. The symptoms were initially triggered spontaneously from daily life, three months prior to his first visit to a local hospital. C-spine MRI revealed CDH and he received physiotherapy for 20 days, three times at the hospital. Then he was hospitalised and received 3 sessions of nerve block injection therapy and traction therapies. The pain remained and he was admitted to Jaseng Hospital of Korean Medicine<sup>36</sup> and received the treatment for 24 days.

At admission, the patient reported a VAS score of 10 and a NDI score of 30. C-spine ROM reported a limitation in left lateral bending of 35° (normal <45°). A positive Spurling test, Roos test and a positive on the left side in Adson's test were also reported. C-spine x-ray (Fig. 5) revealed straightening of the c-spine. The c-spine MRI scan (Fig. 6)




Fig. 3 C-spine x-ray (lateral): disc space narrowing, posterior marginal spurring and degenerative endplate change at C4/5, C5/6 and C6/7.




Fig. 4 Body chart – case 2.

revealed disc herniations at C-2/3, C3/4, C4/5 and C5. At discharge, the patient's VAS score was 2 and the NDI score was 27. Negatives were reported in all three physical examinations (Spurling test, Adson's test, and Roos test).

**Examinations**

There are no definitive guidelines for the diagnosis of TOS and as such, a combination of both physical examination and diagnostic imaging may be



Fig. 5 C-spine x-ray (lateral): straightening in c-spine.




Fig. 6 C-spine MRI: C2/3, C3/4, C4/5 and C5/6 mild central disc protrusion.

required.<sup>18</sup> In the two cases, the physical examination included Spurling's test, Adson's test, and Roos test. All tests were positive in both patients at admission<sup>19</sup> and negative responses at discharge. Spurling's test involves a combination of cervical rotation and flexion with compression and a positive response is generally suggestive of CDH.<sup>2,20</sup> Positive responses with both Adson's test and Roos test are suggestive of TOS.<sup>15,21</sup> Adson's test<sup>2</sup> requires the rotation and extension of the cervical spine towards dysfunctional side while the assessor laterally rotates, abducts and extends the patient's shoulder monitoring the patient's radial pulse. Positive indication was observed with the changes in the pulse. Roos test required the patient to open and close both hands repeatedly with the arms positioned in abduction and extension for 30 s–3 min. Aggravation of the existing symptoms was observed in both cases presented here therefore both tests were considered positive.<sup>2</sup>

#### Outcome measurements

The Visual Analogue Scale (VAS)<sup>22,23</sup> and Neck Disability Index (NDI)<sup>24</sup> were administered both at admission and discharge to measure the degree of pain and evaluate functional changes. The VAS is the most commonly used self-report subjective scale for measuring the degree of pain experienced by a patient.<sup>22–24</sup> VAS scores range from 0 to 10, with 0 being no pain and 10 being the severest.<sup>22</sup> The NDI assesses the impact of cervical spine pain on a number of activities of daily living.<sup>25,26</sup>

#### Treatments

##### Manual therapy

The two cases that are the subject of the current report presented at the Jaseng Hospital in Korea. The hospital provides both western and Korean medical services. Patients are offered a standardised treatment package that consists of manual

therapy, and additional Korean medicine treatments on a regular basis. Upon the diagnosis of a patient's condition, the treatment approach is specifically tailored for the patient.

Both patients received manual therapy sessions every two or three days a week for approximately 20 minutes per session. MET and LAS were performed as the main techniques with additional treatments including herbal medicine, acupuncture and pharmacopuncture. MET and LAS were applied specifically targeting the scalene muscles, ascended first rib and the cervical vertebrae. The following techniques were repeated throughout the entirety of the patient's hospital stay.

##### Muscle energy technique (MET)

MET is a manual therapy technique<sup>16</sup> that has been practiced by many physicians in different clinical fields, and is often used for treating muscle weakness and shortening, restricted joint range of motion, and other related problems.<sup>28</sup> MET requires accuracy and precision in the diagnostic process, and is based on a biomechanical system that assesses restriction of functional movement.<sup>28</sup>

MET for the treatment of neck related symptoms was performed targeting the scalene muscles and it was repeated for 3–5 times at each session until no further increase in range of motion (ROM) were observed.<sup>29</sup>

##### Ligamentous articular strain (LAS)

Ligamentous Articular Strain (LAS) is an osteopathic manipulative technique intended to restore normal functional and physiological movement of the tissues.<sup>17</sup> LAS targets the somatic dysfunction utilising compression or decompression of the joints and fascia to achieve this balance.<sup>24</sup>

To assess and treat the ascended first rib, the outer surface of the patient's cervical vertebrae and the costal surface of the first rib were palpated using the tips of both thumbs. Constant pressure was applied to the first rib until a sense of 'giving way' was achieved.<sup>30</sup>

For the cervical spine, the practitioners hands were placed under the both sides of the patient's base of the skull for support. The middle fingers were removed by approximately 0.5 inches and were placed on the patient's cervical dysfunctional site so that the tips of the fingers remained in contact with the area while moving the operator's fingers to the anterior and superior direction. Finally, the hands were bent slightly to pull the fingers towards the thumbs. Constant pressure was applied on the contact areas of the fingers until full relaxation was achieved.<sup>30</sup>

To address the anterior cervical fascia and the scalene muscle, the tips of both thumbs were placed on the patient's lateral side of supraclavicular fossa and pressed downwards towards the patient's legs. Constant pressure was applied and once the tissue had released, the thumbs were pulled outwards, towards acromioclavicular joint. Due to the sensitivity of this area, only minimal pressure is required to release the muscle. The release of the anterior cervical fascia also contributes to the release of the scalene muscle.<sup>30</sup>

#### Other treatments

##### Herbal medicine

120 ml of Chungpa-jeon (named GCSB-5) in decoction form was prescribed and taken three times daily. Chungpa-jeon is traditionally indicated for spinal disorders for the effects of anti-inflammatory<sup>31</sup> and nerve protection.<sup>32</sup>

##### Acupuncture

Acupuncture treatment was conducted daily on the local area to relieve tension in the muscles, and ligaments in the anterior and posterior neck and shoulder area. The selected acupoints were LI18, LI17, SI17, SI16, TE16, GB20, GB21, LI16, ST12, TE15. LI18 is between the anterior and posterior parts of the sternocleidomastoid muscle (SCM), at the level of the superior border of the thyroid cartilage, and LI17 is directly inferior to LI18, just posterior to the SCM. SI17 is posterior to the mandibular angle, in the depression directly anterior to the SCM, and SI16 is posterior to the SCM, at the level of the superior border of the thyroid cartilage. TE16 is posterior to the mandibular angle, in the depression directly posterior to the SCM. GB20 and GB21 are in the posterior neck and shoulder region; GB20 is inferior to the occiput, and between the origins of the SCM and trapezius, and GB21 is at the midpoint between the C7 spinous process and the lateral end

of the acromion. LI16 is located in the depression between the acromial end of the clavicle and scapular spine, lateral to the suprascapular fossa. ST12 is situated in the depression superior to the clavicle, and TE15. TE15 is in the posterior scapular region, directly superior to the superior scapular angle.<sup>33</sup>

##### Pharmacopuncture

2 cc of Hwangryunhaedok-tang (HRHD-T) pharmacopuncture<sup>34</sup> was injected on the same selected acupuncture points three times daily. HRHD-T pharmacopuncture is used as an anti-inflammatory.<sup>35</sup>

#### Discussion

In both cases that are the subject of the current paper, CDH was confirmed by MRI scans. The common symptom across these two cases was interrupted sleep due to pain and numbness in the arm and hand. Neurological symptoms in CDH are frequently related to the neurological dysfunction around the central spinal structures, whereas in TOS, the symptoms are typically produced by compression of neurovascular structures in the periphery. Furthermore, TOS often causes the pain and numbness to continue throughout the night and day<sup>38</sup> and 'relaxation symptoms' are manifested when the compressed soft tissues are released.<sup>39</sup> Differentiating between CDH and TOS is difficult using the clinical history alone, and there are cases where CDH and TOS have concurrently occurred. Thus, careful consideration of the incidence of both CDH and TOS is required when the clinical symptoms are present in the upper extremities.

The two cases were diagnosed with CDH prior to their admission to Jaseng hospital of Korean Medicine. They both received standard medical treatments for CDH (nerve block therapy, medications, injections etc.) but no improvement was observed. We were able to speculate that TOS could be the cause of the presenting complaints due to the lack of response to the standard medical interventions, rather than identifying TOS according to the patients' clinical signs and symptoms. Orthopaedic examination suggested a diagnosis of TOS, and significant improvement was achieved through application of MET and LAS for the treatment of this condition.

MET and LAS were considered to be effective treatments for these two cases. MET was used to release the shortened scalene muscle to obtain

balance and relieve pain. The LAS<sup>30</sup> was specifically used to release the ascended first rib back and to release anterior cervical fascia. In the first case, a total of 11 sessions of MET and LAS were conducted once every two days for 21 days. In case 2, a total of 12 sessions of MET and LAS were conducted once every 2 days for 24 days. The interventions resulted in significant improvements in the VAS, NDI and in the three orthopaedic tests for TOS.

This report investigates only two successful treatment cases of CDH with TOS. Given this, as well as the use of other interventions, it is difficult to draw conclusions as to the effectiveness of MET and LAS. However the successful outcome presented here suggests that non-surgical approaches for CDH with TOS are worthwhile pursuing, because patients often respond to conservative treatments well enough to achieve significant improvement.<sup>40</sup>

### Conclusion

Surgical approaches in the treatment of spinal disorders, particularly with symptoms in the upper extremities, are frequently observed in many practices. As a result, there are a number of published reports available discussing the surgical indications and approaches for cervical disc herniations, persistent cervical spine pain and persistent neurological changes in the upper extremity. There are however, few reports on the outcomes achieved through manual therapy approaches for the treatment of CDH with TOS. Further research is required into this area and the current case reports suggest that manual therapy may be a useful treatment approach when these conditions occur concurrently.

### Conflict of interest

None declared.

### Ethical approval

None declared.

### Funding

None declared.

### Acknowledgements

This work is supported by Jaseng Medical Foundation.

### References

1. Davidovic LB, Kostic DM, Jakovljevic NS, Kuzmanovic IL, Simic TM. Vascular thoracic outlet syndrome. *World J Surg* 2003;27:545–50.
2. Watson LA, Pizzari T, Balster S. Thoracic outlet syndrome part 1: clinical manifestations, differentiation and treatment pathways. *Man Ther* 2009;14:586–95.
3. Radhakrishnan K, Litchy WJ, O'Fallon WM, Kurland LT. Epidemiology of cervical radiculopathy. A population-based study from Rochester, Minnesota, 1976 through 1990. *Brain* 1994;117:325–35.
4. Carrette S, Fehlings MG. Clinical practice. cervical radiculopathy. *N Engl J Med* 2005;353:392–9.
5. Manchikanti L, Cash KA, Pampati V, Wargo BW, Malla Y. Management of chronic pain of cervical disc herniation and radiculitis with fluoroscopic cervical interlaminar epidural injections. *Int J Med Sci* 2012;9:424–34.
6. Wainner RS, Gill H. Diagnosis and nonoperative management of cervical radiculopathy. *J Orthop Sports Phys Ther* 2000;30:728–44.
7. Roos DB. The place for scalenectomy and first-rib resection in thoracic outlet syndrome. *Surgery* 1982;92:1077–85.
8. Samarasam I, Sadhu D, Agarwal S, Nayak S. Surgical management of thoracic outlet syndrome: a 10-year experience. *ANZ J Surg* 2004;74:450–4.
9. Yanaka K, Asakawa H, Matsumaru Y, Kujiraoka Y, Nose T. Diagnosis of vascular compression at the thoracic outlet using magnetic resonance angiography. *Eur Neurol* 2004;51:122–3.
10. Lindgren KA, Oksala I. Long-term outcome of surgery for thoracic outlet syndrome. *Am J Surg* 1995;169:358–60.
11. Atasoy E. Thoracic outlet compression syndrome. *Orthop Clin North Am* 1996;27:265–303.
12. Rayan GM. Thoracic outlet syndrome. *J Shoulder Elbow Surg* 1998;7:440–51.
13. Cooke RA. Thoracic outlet syndrome – aspects of diagnosis in the differential diagnosis of hand-arm vibration syndrome. *Occup Med (Lond)* 2003;53:331–6.
14. Brooke BS, Freischlag JA. Contemporary management of thoracic outlet syndrome. *Curr Opin Cardiol* 2010;25:535–40.
15. Sanders RJ, Hammond SL, Rao NM. Diagnosis of thoracic outlet syndrome. *J Vasc Surg* 2007;46:601–4.
16. DeStefano LA. *Greenman's principles of manual medicine*. United States: Lippincott Williams & Wilkins, a Wolters Kluwer Business; 2011.
17. DiGiovanna EL, Schiowitz S, Dowling DJ. Ligamentous articular strain technique and balanced ligamer. In: Somers D, editor. *An osteopathic approach to diagnosis and treatment*. USA: Lippincott Williams & Wilkins; 1991. p. 103–6.
18. Nichols D, Selger C. Diagnosis and treatment of a patient with bilateral thoracic outlet syndrome secondary to anterior subluxation of bilateral sternoclavicular joints: a case report. *Physiother Theory Pract* 2013;29:562–71.
19. Novak CB, Collins ED, Mackinnon SE. Outcome following conservative management of thoracic outlet syndrome. *J Hand Surg Am* 1995;20:542–8.

20. Anekstein Y, Blecher R, Smorgick Y, Mirovsky Y. What is the best way to apply the spurling test for cervical radiculopathy? *Clin Orthop Relat Res* 2012;470:2566–72.
21. Sanders RJ, Hammond SL, Rao NM. Thoracic outlet syndrome: a review. *Neurologist* 2008;14:365–73.
22. McCormack HM, Horne DJ, Sheather S. Clinical applications of visual analogue scales: a critical review. *Psychol Med* 1988;18:1007–19.
23. Tiplady B, Jackson SH, Maskrey VM, Swift CG. Validity and sensitivity of visual analogue scales in young and older healthy subjects. *Age Ageing* 1998;27:63–6.
24. Mehta S, Macdermid JC, Carlesso LC, McPhee C. Concurrent validation of the DASH and the QuickDASH in comparison to neck-specific scales in patients with neck pain. *Spine (Phila Pa 1976)* 2010;35:2150–6.
25. Pietrobon R, Coeytaux RR, Carey TS, Richardson WJ, DeVellis RF. Standard scales for measurement of functional outcome for cervical pain or dysfunction: a systematic review. *Spine (Phila Pa 1976)* 2002;27:515–22.
26. Vernon H. The neck disability index: state-of-the-art, 1991–2008. *J Manipulative Physiol Ther* 2008;31:491–502.
28. Mitchell JF, Mitchell P. *The MET manual*, vol. 1. East Lansing, Michigan: MET Press; 1995.
29. Chaitow L. *Muscle Energy techniques with DVD-ROM*. UK: Churchill Livingstone; 2006.
30. Speece AC, Conrad TW, Simmons LS. *Ligamentous articular strain: osteopathic manipulative techniques for the body*. Revised ed. US: Eastland Press; 2009.
31. Chung HJ, Lee HS, Shin JS, Lee SH, Park BM, Youn YS, et al. Modulation of acute and chronic inflammatory processes by a traditional medicine preparation GCSB-5 both in vitro and in vivo animal models. *J Ethnopharmacol* 2010;130:450–9.
32. Kim TH, Yoon SJ, Lee WC, Kim JK, Shin J, Lee S, et al. Protective effect of GCSB-5, an herbal preparation, against peripheral nerve injury in rats. *J Ethnopharmacol* 2011;136:297–304.
33. World Health Organization, Regional Office for the Western Pacific. *WHO standard acupuncture point locations in the Western Pacific Region*. WHO Regional Office for the Western Pacific; 2008.
34. Korean Pharmacopuncture Institute. *Pharmacopunctureology: principles and clinical application*. Republic of Korea: Elsevier Korea LLC; 2011.
35. Kim NK, Lee DH, Seo HS, Sun SH, Oh YL, Kim JE, et al. Hwangryunhaedoktang in adult patients with atopic dermatitis: a randomised, double-blind, placebo-controlled, two-centre trial – study protocol. *BMC Complement Altern Med* 2011;11: 68-6882-11-68.
36. Stevens L, Duarte H, Park J. Promising implications for integrative medicine for back pain: a profile of a Korean hospital. *J Altern Complement Med* 2007;13:481–4.
38. Novak CB. Conservative management of thoracic outlet syndrome. *Semin Thorac Cardiovasc Surg* 1996;8:201–7.
39. Yao STJ, Pearce HW. *Vascular surgery-therapeutic strategies*. USA: Pmph USA; 2010.
40. Saal JS, Saal JA, Yurth EF. Nonoperative management of hemiated cervical intervertebral disc with radiculopathy. *Spine (Phila Pa 1976)* 1996;21:1877–83.

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

**ScienceDirect**